

For answers, send email to: admin@tutor-homework.com.

Include file name: Chemistry_Worksheet_0107

Price: \$3

(c) 2012 www.tutor-homework.com: Tutoring, homework help, help with online classes.

1. Compression Work1

Calculate the quantity of work, in J, when a gas is compressed from 2.3 L to 0.6 L by a constant external pressure of 7.79 atm.

2. First Law (-w,+q) find dU1

Determine the change in internal energy (in kJ) of a system if the system does 239 kJ of work on the surroundings and 235 kJ of heat is transferred to the system from the surroundings?

3. Expt Det of Metal Specific Heat

When a 149 g sample of a metal at 241°C is dropped in 314 g of water, the temperature of the water increases from 25°C to 30°C. Given that the specific heat of water is 4.184 J/g°, what is the heat capacity of the metal (in J/g°).

4. The reaction of glucose C₆H₁₂O₆ with oxygen gas p -

The reaction of glucose (C₆H₁₂O₆) with oxygen gas produces carbon dioxide and water. The energy released during this reaction is 2803 kJ/mol of glucose. If 18 moles of oxygen are used, how many kilocalories are released during the reaction?

Student
Response

a. 2.010×10^3

b. 6695

c. 2003

d. 12.69

e. 5987

5. - A reaction took place in four steps. Each step gave -

A reaction took place in four steps. Each step gave off 100 cal of heat. In the fourth step there was also a decrease in volume by 76.0 L at a pressure of 800 mm Hg. What was the energy change in joules for the overall reaction?

Student
Response

- a. +228
- b. -1.59×10^3
- c. -6.12×10^4
- d. -480
- e. $+6.43 \times 10^3$

6. Chapter05-047


Given the following reactions

is _____ kJ.

Student Response
a. 145.7
b. 343.9
c. -343.9
d. 17.5
e. -145.7

7. Enthalpy of Reaction from Formation Values Calc.

Calculate the enthalpy change ΔH_{rxn} (in kJ) for the following hypothetical reaction:



The formation enthalpies are tabulated as follows:

	A	B	C	D
ΔH_f° (kJ/mol)	329.5	-169.6	201.3	-169.7

8. Bomb Calorimeter Calc

The heat capacity of a bomb calorimeter was determined by burning 1.3 moles of compound X (heat of combustion = 575 kJ/mol) in the bomb. The temperature increased by 32° . When 0.8 moles of compound Y is combusted in the same calorimeter, the temperature of the calorimeter increases by 22.5° . What is the heat

capacity of compound Y?

9-20 KINETICS

9. - For the reaction $2\text{SO}_3 \rightarrow 2\text{SO}_2 + \text{O}_2$ calculate the aver -

For the reaction $2\text{SO}_3 \rightarrow 2\text{SO}_2 + \text{O}_2$ calculate the average rate, $\Delta[\text{SO}_3]/\Delta t$, between 10.0 min and 40.0 min for the loss of SO_3 .

Time, min:	0.0	10.0,	20.0	30.0,	40.0	50.0
$[\text{SO}_3]$, M	0.124	0.092	0.068	0.050	0.037	0.028
$[\text{SO}_2]$, M:	0.0	0.032	0.056	0.074	0.087	0.096
$[\text{O}_2]$, M	0.0	0.016	0.028	0.037	0.044	0.048

Student Response

a. $2.2 \times 10^{-4} \text{ M min}^{-1}$

b. $1.8 \times 10^{-3} \text{ M min}^{-1}$

c. $1.1 \times 10^{-3} \text{ M min}^{-1}$

d. $5.5 \times 10^2 \text{ M min}^{-1}$

e. $4.6 \times 10^2 \text{ M min}^{-1}$

10. Determine Rate (monitor reactant B)

The reaction



is monitored by measuring the decrease in the concentration of B over time. The concentration of B was found to be 14.3 M after 29.2 minutes had elapsed, and 8.78 M after 122.2 minutes had elapsed. What is the reaction rate (in M/min)?

11. Exp Rate Law (A+B Rx)

For a reaction in which A and B react to form C, the following initial rate data were obtained:

Experiment	[A], M	[B], M	initial rate, M/s:
1	0.2	0.2	0.50
2	0.4	0.2	2.00
3	0.8	0.2	8.00
4	0.2	0.4	1.00
5	0.2	0.8	2.00

The rate law for this reaction has the general form

$$\text{Rate} = k[\text{A}]^a[\text{B}]^b$$

Determine (1) a, the order with respect to A; (2) b, the order with respect to B; (3) the overall order

12. Value of K

For the hypothetical reaction $\text{A} + \text{B} \rightarrow \text{products}$, the rate law is found to be

$$R = k[\text{A}]^1[\text{B}]^2$$

when $[\text{A}]_0 = 5.9 \text{ M}$ and $[\text{B}]_0 = 0.7 \text{ M}$, the initial rate = $3.23 \times 10^2 \text{ M s}^{-1}$. What is the value of the rate coefficient, k ?

13. - In a first-order rate expression, what are the unit -

In a first-order rate expression, what are the units for the rate constant?

Student Response
a. time^{-1}
b. time
c. $(\text{M time})^{-1}$
d. time M^{-1}
e. unitless

14. t-Half for 1 order

Determine the half-life (in s) for the reaction $\text{A} \rightarrow \text{products}$, given that the reaction is first order, $k = 1.707 \text{ s}^{-1}$ and $[\text{A}]_0 = 4.16759$

15. First Order A given t1/2 and t


A certain first order process



has a half-life of 41.9 minutes. If a reaction vessel is initially charged with 445 mmHg of A, what will the partial pressure of A be after 25.8 minutes have elapsed.

16. Collision Attributes

Consider the reaction




1. In order for a collision between H_2 and I_2 to lead to products, what two attributes must the collision possess?
2. Explain how each of the following changes would improve the odds of a successful reaction event
 - a. Increasing temperature
 - b. Increasing the concentrations of H_2 and/or I_2
 - c. Adding a catalyst

17. Endo Profile

Consider the following energy profile diagram for the reaction



1. Is this an exothermic or endothermic reaction?
2. What is point X called?
3. ΔH_{rxn} is _____ kJ/mol
4. The activation energy for the reaction $\text{A} \rightarrow \text{B}$ is _____ kJ/mol
5. The activation energy for the reaction $\text{B} \rightarrow \text{A}$ is _____ kJ/mol



18. Arrhenius Calc Find Ea

The hypothetical reaction



has a rate coefficient of $3.0 \times 10^{-15} \text{ s}^{-1}$ at 187 K and a rate coefficient of $1.6 \times 10^{-7} \text{ s}^{-1}$ at 875 K. What is the activation energy for this reaction (in kJ/mol). $R = 8.314$

J/(molK)

19. Molecularity 3


Assign the correct molecularity to each of the following elementary reactions:

1. $2\text{NO}_2 \rightarrow 2\text{NO} + \text{O}_2$
2. $2\text{NO} + \text{Br}_2 \rightarrow 2\text{NOBr}$
3. $\text{SO}_2\text{Cl}_2 \rightarrow \text{SO}_2 + \text{Cl}_2$
4. $\text{NO}_2 + \text{F}_2 \rightarrow \text{NO}_2\text{F} + \text{F}$

Unimolecular
Termolecular
Bimolecular

20. Steady-State Mechanism Ozone

A mechanism for the decomposition of ozone: $2\text{O}_3(\text{g}) \rightarrow 3\text{O}_2$ is:



Applying the steady-state approximation the following rate law is obtained.

$$R = \frac{k'X^2}{X + k''Y}$$

What is X and Y?

X
Y

- a. O_3
- b. O_2
- c. O